

# BJP Psych

The British Journal of Psychiatry

**Assessing early-onset hallucinations in 'the touch-screen generation'**

Morgane Demeulemeester *et al*

**Letters, green cards, telephone calls and postcards: systematic and meta-analytic review of brief contact interventions for reducing self-harm, suicide attempts and suicide**

Allison Milner *et al*

**Adverse childhood events and psychosis in bipolar affective disorder**

Rachel Upthegrove *et al*

**Monoamine oxidase A gene promoter methylation and transcriptional downregulation in an offender population with antisocial personality disorder**

Dave Checknita *et al*



- A9 Editorial Board  
A11 Highlights of this issue

## Editorials

- 177 **An integrated biopsychosocial model of childhood maltreatment and psychosis**  
V. Barker, A. Gumley, M. Schwannauer and S. M. Lawrie

- 180 **You're not welcome in here – poems by doctors**  
Hinesh Topiwala

- 181 **Assessing early-onset hallucinations in the touch-screen generation**  
M. Demeulemeester, F. Kochman, B. Fligans, A. J. Tabet, P. Thomas and R. Jardri

## Review article

- 184 **Letters, green cards, telephone calls and postcards: systematic and meta-analytic review of brief contact interventions for reducing self-harm, suicide attempts and suicide**  
A. J. Milner, G. Carter, J. Pirkis, J. Robinson and M. J. Spittal

## Papers

- 191 **Adverse childhood events and psychosis in bipolar affective disorder**  
R. Upthegrove, C. Chard, L. Jones, K. Gordon-Smith, L. Forty, I. Jones and N. Craddock
- 198 **Neural responses to dynamic multimodal stimuli and pathology-specific impairments of social cognition in schizophrenia and depression**  
C. Regenbogen, T. Kellermann, J. Seubert, D. A. Schneider, R. E. Gur, B. Derntl, F. Schneider and U. Habel
- 206 **Common and disorder-specific neural responses to emotional faces in generalised anxiety, social anxiety and panic disorders**  
G. A. Fonzo, H. J. Ramsawh, T. M. Flagan, S. G. Sullivan, A. Letamendi, A. N. Simmons, M. P. Paulus and M. B. Stein
- 216 **Monoamine oxidase A gene promoter methylation and transcriptional downregulation in an offender population with antisocial personality disorder**  
D. Checknita, G. Maussion, B. Labonté, S. Comaj, R. E. Tremblay, F. Vitaro, N. Turecki, A. Bertazzo, G. Gobbi, G. Côté and G. Turecki

- 222 **Out-of-body experiences – in 100 words**  
Peter Fenwick

- 223 **Droperidol v. haloperidol for sedation of aggressive behaviour in acute mental health: randomised controlled trial**  
L. Calver, V. Drinkwater, R. Gupta, C. B. Page and G. K. Isbister

- 229 **The ACCESS study: Zelen randomised controlled trial of a package of care for people presenting to hospital after self-harm**

S. Hatcher, C. Sharon, A. House, N. Collins, S. Collings and A. Pillai

- 237 **Factors associated with inconsistency in self-reported mild traumatic brain injury over time among military personnel in Iraq**

N. W. Nelson, C. R. Anderson, P. Thuras, S. M. Kehle-Forbes, P. A. Arbisi, C. R. Erbes and M. A. Polusny

- 244 **Locked in – extra**  
Andrew Gardner

- 245 **Latent structure of the proposed ICD-11 post-traumatic stress disorder symptoms: implications for the diagnostic algorithm**

D. Forbes, E. Lockwood, M. Creamer, R. A. Bryant, A. C. McFarlane, D. Silove, A. Nickerson and M. O'Donnell

- 251 **Gertrude Parker: a life in an asylum – 100 years ago**  
Joy Parker

## Columns

- 252 Correspondence  
255 Book reviews  
257 Contents of *BJPsych Advances*  
258 Contents of the *American Journal of Psychiatry*  
259 Kaleidoscope  
261 From the Editor's desk

## Cover picture

*Spirited* (2013).  
Megan Winton (b. 1997)

Megan writes: 'This piece was created to show the four primates, Jethro (white-faced Saki monkey) and spider monkeys Mickory, Flint and Pumpkin. All four of these primates were re-homed at Monkey World ape and monkey rescue centre last year, as their owners had to move and asked Monkey World to take them in. The previous owners had great experience in caring for these primates and had raised them from infancy, so it was difficult to see them go. I wanted to express their lovely personalities and create a spellbinding atmosphere, as this is how these monkeys made me feel. The fantasy element to the painting helps to give a more serene aspect, which is how I like to escape from real life and my mental health issues.

'My problems are OCD, PTSD and selective mutism. I have been unwell for 5 years, so have lost 5 years of my life. I have recently felt as if I'm not really 17 years old. However, I now have more control over my OCD symptoms, although the mutism is still a problem at times of anxiety and stress'.

We are always looking for interesting and visually appealing images for the cover of the *Journal* and would welcome suggestions or pictures, which should be sent to Dr Allan Beveridge, British Journal of Psychiatry, 21 Prescot Street, London E1 8BB, UK or [bjp@rcpsych.ac.uk](mailto:bjp@rcpsych.ac.uk).



## Highlights of this issue

By Sukhwinder S. Shergill

### PTSD and aggression

The new International Classification of Diseases version 11 (ICD-11) is awaited, and the contentious diagnosis of post-traumatic stress disorder (PTSD) has been simplified into three factors representing symptoms related to re-experiencing, avoidance and hyperarousal. However, Forbes and colleagues (pp. 245–251) found that a two-factor solution collapsing re-experiencing and avoidance factors provided an equivalent fit to the three-factor model. They suggest that the two-factor model, derived from their factor analysis on symptoms present in 613 patients with PTSD, provides a more useful inclusive algorithm for diagnosis. The management of acute aggression or behavioural disturbance is always difficult to assess in clinical studies; Calver and colleagues (pp. 223–228) are to be commended for their randomised controlled study of intramuscular droperidol *v.* haloperidol. They found that both resulted in effective sedation within 20–25 minutes, with no differences in efficacy between them; however, there were some trends towards greater use of additional sedation with haloperidol, but with fewer adverse effects.

### Childhood adversity and psychosis

There is a reported association between childhood adversity and the development of psychosis. Upthegrove *et al* (pp. 191–197) did not confirm this link in their large study of patients with bipolar disorder. However, there were specific associations of hallucinations with childhood sexual abuse, death of loved one and being victimised. These were not associated with the development of delusions. The authors suggest that, given the phenomenological specificity of their findings, future studies need to examine these subtypes in more detail if we are to identify and improve outcomes in those at increased risk. On a related topic, Barker and colleagues (pp. 177–180) review putative biological mechanisms that could underpin the association between childhood adversity and psychosis, with a focus on alterations within the hypothalamic–pituitary–adrenal axis mediated by expression of brain-derived neurotrophic factor in the hippocampus and the neuropeptide oxytocin. In their editorial, they propose epigenetic mechanisms as the key feature by which environmental effects can influence longer-lasting phenotypic change, and use attachment theory as a psychological construct to frame these changes at the cognitive level. They caution against a simplistic translation into therapy, highlighting the caveats that

much of this work is in animal models, with weaker data in humans. The use of technology to improve assessment of hallucinatory symptoms is the focus of an editorial by Demeulemeester *et al* (pp. 181–183). They review the increasing use of health apps in assessing symptoms in mental health; and then describe the development of the Multisensory Hallucination Scale for Children (MHASC), designed to be used on tablets, and the advantages of using an approach that is set as part game and part standardised test.

### Reducing the incidence of repeated self-harm

There is an enormous personal, economic and societal cost associated with suicide and self-harm. Given the difficulty in predicting and preventing self-harm *de novo*, it is essential to optimise our response to people presenting with self-harm, in order to reduce the frequency of any subsequent attempts. In a review of brief contact interventions after self-harm Milner and colleagues (pp. 184–190) found that overall these methods did not confer a significant benefit on repetition rates or suicide. The authors suggest that larger-scale studies are necessary to examine some of the more promising individual approaches, such as postcard interventions. A study of a package of measures, including postcards and problem-solving therapy, in people presenting to hospital after self-harm supported the findings of the review; there were no significant differences in outcomes at a 1 year follow-up post-intervention. Hatcher *et al* (pp. 229–236) suggest that despite the lack of consistent positive differences, low-cost interventions such as postcards may still be worth implementing as they may be helpful for some people.

### Neural change in psychiatric disorder

It is unusual to compare neural signatures across many different diagnostic groups; Fonzo and colleagues (pp. 206–215) demonstrate that amygdala activation was related to greater negative affect or anxiety, whereas patients with panic disorder were more likely to show activation in the posterior insula. The authors suggest that this multidagnostic approach has utility in clarifying dimensional characterisation of mental illness. A similar approach was used by Regenbogen *et al* (pp. 198–205), who compared brain activation using functional magnetic resonance imaging during a dynamic social cognition task in patients with schizophrenia and patients with depressive illness. They suggest that patients with schizophrenia were characterised by disturbance of neural networks associated with processing semantic context, whereas patients with depression showed greater change in mentalising networks. They suggest that dynamic social stimuli may offer useful probes for differentiating between different psychiatric diagnoses.