


# BJPsych

The British Journal of Psychiatry



**Does the stigma  
of mental illness  
contribute to  
suicidality?**

Nicolas Rüsch *et al*

**Psychiatric  
resilience:  
longitudinal twin  
study**

Ananda Amstadter *et al*

**Diffusion tensor  
imaging predictors  
of treatment  
outcomes in major  
depressive disorder**

Mayuresh Korgaonkar  
*et al*

**Clinical and social  
factors associated  
with attention-  
deficit hyperactivity  
disorder medication  
use:  
population-based  
longitudinal study**

Cédric Galéra *et al*



# BJPsych

## Contents

- A13 Editorial Board  
A15 Highlights of this issue

### Editorial

- 257 **Does the stigma of mental illness contribute to suicidality?**  
N. Rüsch, A. Zlati, G. Black and G. Thornicroft

- 259 **Intense romantic love in a 12-century Persian medical encyclopaedia – extra**  
Behnam Dalfardi, Golnoush Sadat Mahmoudi Nezhad and Ahmad Ghanizadeh

### Review articles

- 260 **Excess of non-right-handedness in schizophrenia: meta-analysis of gender effects and potential biases in handedness assessment**  
M. Hirnstein and K. Hugdahl
- 268 **Psychotherapy for subclinical depression: meta-analysis**  
P. Cuijpers, S. L. Koole, A. van Dijke, M. Roca, J. Li and C. F. Reynolds III

- 274 **The Man He Killed – poem**  
Thomas Hardy

### Papers

- 275 **Psychiatric resilience: longitudinal twin study**  
A. B. Amstadter, J. M. Myers and K. S. Kendler
- 281 **Invited commentary on . . . Psychiatric resilience: longitudinal twin study**  
J. Wertz and C. M. Pariante
- 283 **Suicide rates in children aged 10–14 years worldwide: changes in the past two decades**  
K. Kölves and D. De Leo
- 286 **Childhood family income, adolescent violent criminality and substance misuse: quasi-experimental total population study**  
A. Sariaslan, H. Larsson, B. D'Onofrio, N. Långström and P. Lichtenstein
- 291 **Clinical and social factors associated with attention-deficit hyperactivity disorder medication use: population-based longitudinal study**  
C. Galéra, J.-B. Pingault, G. Michel, M.-P. Bouvard, M. Melchior, B. Falissard, M. Boivin, R. E. Tremblay and S. M. Côté

- 297 **Postpartum psychosis – in 100 words**  
Roch Cantwell

- 298 **Pathways of care of women in secure hospitals: which women go where and why**  
A. Bartlett, N. Somers, M. Fiander and M. A. Harty

- 307 **Internet-delivered exposure-based cognitive-behavioural therapy and behavioural stress management for severe health anxiety: randomised controlled trial**  
E. Hedman, E. Axelsson, A. Göring, C. Ritzman, M. Ronnheden, S. El Alaoui, E. Andersson, M. Lekander and B. Ljótsson

- 315 **White matter integrity and late-life depression in community-dwelling individuals: diffusion tensor imaging study using tract-based spatial statistics**  
S. Reppermund, L. Zhuang, W. Wen, M. J. Slavin, J. N. Trollor, H. Brodaty and P. S. Sachdev

- 321 **Diffusion tensor imaging predictors of treatment outcomes in major depressive disorder**  
M. S. Korgaonkar, L. M. Williams, Y. J. Song, T. Usherwood and S. M. Grieve

- 328 **A dissection of 'Paranoid Android' – psychiatry in music**  
Parashar Ramanuj

### Columns

- 329 Correspondence  
330 Correction  
331 Book reviews  
332 Contents of *Advances in Psychiatric Treatment*  
333 Contents of the *American Journal of Psychiatry*  
334 Kaleidoscope  
336 From the Editor's desk

### Cover picture


*The Demonstration* (late 1950s/early 60s, poster paint on paper). Mary Bishop (b: currently unknown; d: 1990).

Mary Bishop's father had died in the First World War. She became psychotic while serving in the Women's Auxiliary Air Force during the Second World War. She was admitted to Netherne Hospital around 1946 and discharged circa 1983. During these years, she painted thousands of pictures in Edward Adamson's Netherne art studios depicting her isolation, depression, anger, and despair. About 300 works survive and are part of the recently found Adamson Collection. Adamson was a British artist who pioneered the use of art as therapy. In his book *Art as Healing* (1984; re-printed 2014, Adamson Collection Trust) he comments on *The Demonstration*: 'Here she illustrates exactly how it feels to be demonstrated in front of a group of medical students. She feels that she is entirely naked before an audience of blockheads' (p.21). Adamson adds: 'She told me that her existence was only made bearable by the fact that she could come and paint whenever she liked'.

David O'Flynn, Consultant Psychiatrist, SLAM, and Chair, Adamson Collection Trust.

Courtesy of the Adamson Collection Trust.

We are always looking for interesting and visually appealing images for the cover of the *Journal* and would welcome suggestions or pictures, which should be sent to Dr Allan Beveridge, British Journal of Psychiatry, 21 Prescot Street, London E1 8BB, UK or [bjp@rcpsych.ac.uk](mailto:bjp@rcpsych.ac.uk).





## Highlights of this issue

By Kimberlie Dean

**Biological markers and psychotherapeutic interventions for depression**

Two papers in the *BJPsych* this month investigate brain imaging markers in relation to depression outcomes. On the basis that late-life depression has previously been found to be associated with white matter changes on region of interest brain imaging, Reppermund *et al* (pp. 315–320) utilised diffusion tensor imaging and tract-based spatial statistics to examine white matter integrity in a sample of older adults, twice over a 2-year period. Those with current depression had widespread integrity reduction compared with those without depression, with abnormalities particularly apparent in the frontal lobe, association and projection fibres. The authors also found that white matter integrity reduction in a number of specific brain regions predicted later depression at follow-up, indicating potential as a biomarker for such an outcome. The ability of diffusion tensor imaging findings to predict outcome in depression was also examined by Korgaonkar *et al* (pp. 321–328), this time in the context of predicting treatment outcomes in a sample of individuals with major depressive disorder being treated in an out-patient setting. Taking demographic and clinical measures into account, altered connectivity in the cingulum part of the cingulate and stria terminalis tracts predicted remission for those receiving an 8-week course of antidepressant treatment.

Although subclinical depression is common and known to be associated with a range of adverse outcomes, clinical decisions about appropriate treatment are hampered by uncertainty and controversy regarding the evidence for effectiveness. Cuijpers *et al* (pp. 268–274) conducted a meta-analysis of 18 controlled trials of psychological treatment for subclinical depression. They found evidence for a small to moderate effect of psychotherapy on depressive symptoms and a significant impact on incidence of major depression. The authors found the studies to be limited in quality and to vary considerably with regard to characteristics such as the nature of the sample and therapy. The authors noted that the effects found were significantly smaller than those found for psychotherapy when used to treat major depressive disorder.

**Longitudinal studies – of both psychiatric resilience and predictors of poor outcome**

Variability in response to stressful life events is well established but is poorly understood, particularly the mechanisms underlying resilience to such events. Utilising a longitudinal twin study design, Amstadter *et al* (pp. 275–280) found that the difference between actual level of internalising symptoms and that predicted

on the basis of the occurrence of life events (resilience) was underpinned by both genetic and enduring environmental influences, with the effects being roughly equal in impact. In a commentary on the paper, Wertz & Pariente (pp. 281–282) discuss the extent to which the findings could inform development of interventions to enhance resilience after exposure to stressful life events and call for future research to consider a broader range of resilience domains on the basis that, for example, individuals may be resilient to developing psychiatric ill health in response to adversity but not to the development of physical ill health.

Two papers in the *BJPsych* this month present findings from longitudinal studies examining population-level predictors of mental health outcome. Using a quasi-experimental family-based study, Sariaslan *et al* (pp. 286–290) found that childhood family income was not associated with criminality or substance misuse in adolescence once adjustment had been made for unobserved family risk factors. The authors argue that claims of causality based on studies of such associations using traditional epidemiological methods with simple adjustment for covariates should be made with caution. The lack of evidence for causality found when a quasi-experimental family-based design was utilised implies that interventions designed solely to address childhood family income are unlikely to have the desired impact on later risk of criminality or substance misuse. Galéra *et al* (pp. 291–297) examined predictors of attention-deficit hyperactivity disorder (ADHD) medication use in a population-based birth cohort. In addition to ADHD symptoms, a number of social variables, including low maternal education, were also predictive of ADHD medication. Independent of ADHD symptoms, psychiatric comorbidity, including oppositional features beyond those explained by ADHD status, did not predict medication use, but boys were more likely to receive medication than girls. The authors also comment on the finding that parental immigrant status predicted lower medication use.

**Suicide in childhood and in relation to the stigma of mental illness**

Addressing the limited research on suicide patterns in children, Kölves & De Leo (pp. 283–285) obtained data from the World Health Organization Mortality Database and the World Bank for 81 countries and territories for those in the population aged 10–14 years. Over the past two decades, the suicide rate for boys in this age group declined slightly (1.61 per 100 000 to 1.52) but for girls the rate increased slightly (0.85 to 0.94). Regional differences were identified and the authors comment on the potential impact of changes in the quality of data from particular regions over the period of observation. In an editorial focused on the impact of the stigma of mental illness on suicidality, Rüsch *et al* (pp. 257–259) argue that more research is needed to understand what is likely to be an important association between stigma and suicidality, one that, significantly, is potentially modifiable.