

BJPpsych

The British Journal of Psychiatry

Ending lethal discrimination against people with serious mental illness

Holly Taggart
and Sue Bailey

Case study of posts before and after a suicide on a Swedish internet forum

Michael Westerlund
et al

Different outcomes of never-treated and treated patients with schizophrenia: 14-year follow-up study in rural China

Mao-Sheng Ran *et al*

Investigating trajectories of social recovery in individuals with first-episode psychosis: a latent class growth analysis

Jo Hodgekins *et al*



BJPsych

Contents

- A21 Editorial Board
A23 Highlights of this issue

Editorials

- 469 **Ending lethal discrimination against people with serious mental illness**
H. Taggart and S. Bailey
- 471 **Health inequalities and psychosis: time for action**
D. Shiers, T. Bradshaw and J. Campion
- 474 **Out of sight, out of mind**
C. G. D. Brooker and M. Brown

Special article

- 476 **Case study of posts before and after a suicide on a Swedish internet forum**
M. Westerlund, G. Hadlaczky and D. Wasserman

- 482 **Psalm 102: A prayer of the afflicted – psychiatry in the Bible**
Seri Abraham

Review article

- 483 **Mental state decoding impairment in major depression and borderline personality disorder: meta-analysis**
M. J. Richman and Z. Unoka

- 489 ***De praestigiis daemonum*: the origins of psychiatric history-taking – psychiatry in history**
George Huntington

Papers

- 490 **Loci with genome-wide associations with schizophrenia in the Han Chinese population**
Z. Li, Y. Xiang, J. Chen, Q. Li, J. Shen, Y. Liu, W. Li, Q. Xing, Q. Wang, L. Wang, G. Feng, L. He, X. Zhao and Y. Shi
- 495 **Different outcomes of never-treated and treated patients with schizophrenia: 14-year follow-up study in rural China**
M.-S. Ran, X. Weng, C. L.-W. Chan, E. Y.-H. Chen, C.-P. Tang, F.-R. Lin, W.-J. Mao, S.-H. Hu, Y.-Q. Huang and M.-Z. Xiang
- 501 **Accuracy of proactive case finding for mental disorders by community informants in Nepal**
M. J. D. Jordans, B. A. Kohrt, N. P. Luitel, I. H. Komproe and C. Lund
- 507 **Cross-national variations in reported discrimination among people treated for major depression worldwide: the ASPEN/INDIGO international study**
A. Lasalvia, T. Van Bortel, C. Bonetto, G. Jayaram, J. van Weeghel, S. Zoppei, L. Knifton, N. Quinn, K. Wahlbeck, D. Cristofalo, M. Lanfredi, N. Sartorius, G. Thornicroft and the ASPEN/INDIGO Study Group
- 515 **Victimisation in adults with severe mental illness: prevalence and risk factors**
L. D. de Mooij, M. Kikkert, N. M. Lommerse, J. Peen, S. C. Meijwaard, J. Theunissen, P. W. R. A. Duurkoop, A. E. Goudriaan, H. L. Van, A. T. F. Beekman and J. J. M. Dekker
- 523 **Criminal justice pathways to psychiatric care for psychosis**
K. Bhui, S. Ullrich, C. Kallis and J. W. Coid

- 529 **Offerings – poems by doctors**
Cate Bailey

- 530 **Ethnic inequalities in the use of secondary and tertiary mental health services among patients with obsessive-compulsive disorder**

L. Fernández de la Cruz, M. Llorens, A. Jassi, G. Krebs, P. Vidal-Ribas, J. Radua, S. L. Hatch, D. Bhugra, I. Heyman, B. Clark and D. Mataix-Cols

- 536 **Investigating trajectories of social recovery in individuals with first-episode psychosis: a latent class growth analysis**
J. Hodgekins, M. Birchwood, R. Christopher, M. Marshall, S. Coker, L. Everard, H. Lester, P. Jones, T. Amos, S. Singh, V. Sharma, N. Freemantle and D. Fowler

- 543 **Verbatim – in 100 words**
David Brunskill

- 544 **Development of the REFOCUS intervention to increase mental health team support for personal recovery**
M. Slade, V. Bird, C. Le Boutillier, M. Farkas, B. Grey, J. Larsen, M. Leamy, L. Oades and J. Williams

- 551 **Psychometric evaluation of the Questionnaire about the Process of Recovery (QPR)**
J. Williams, M. Leamy, F. Pesola, V. Bird, C. Le Boutillier and M. Slade

Short report

- 556 **Acute intermittent porphyria: comorbidity and shared familial risks with schizophrenia and bipolar disorder in Sweden**
M. Cederlöf, S. E. Bergen, H. Larsson, M. Landén and P. Lichtenstein
- 558 **Postnatal paternal depressive symptoms associated with fathers' subsequent parenting: findings from the Millennium Cohort Study**
S. Nath, G. Russell, T. Ford, W. Kuyken and L. Psychogiou

Columns

- 560 Correspondence
- 563 Contents of *BJPsych Advances*
- 564 Contents of the *American Journal of Psychiatry*
- 565 Kaleidoscope
- 567 From the Editor's desk

Cover picture

Castles (2014).
By Jamie Kamal (b. 1992)

Jamie Kamal is an artist who enjoys photographing what he sees when out walking in the countryside and in the urban environment. Often these photographs are of buildings and views across the city of Edinburgh where he lives. From this starting point, he creates unique images emphasising the shapes, patterns and order of architecture. His signature style features fine lines and intricate dots.

Jamie has severe autism and he is largely non-verbal. His artwork enables him to explore an enjoyable, personal view of his world and to communicate without words. He regularly attends classes with Heather Lucchesi, whose social enterprise organisation, Light Bulb Arts, provides inclusive art experiences for people of all ages and abilities. This summer Jamie's work was on show at The Gallery on the Corner in Edinburgh's New Town and his illustrations appeared in the brochure for the Edinburgh Art Festival Explorers guide.

With thanks to Heather Lucchesi for this submission.

We are always looking for interesting and visually appealing images for the cover of the *Journal* and would welcome suggestions or pictures, which should be sent to Dr Allan Beveridge, British Journal of Psychiatry, 21 Prescott Street, London E1 8BB, UK or bjp@rcpsych.ac.uk.



Highlights of this issue

By Kimberlie Dean

The impact of inequality and discrimination on mental health outcomes

While the prevalence of obsessive-compulsive disorder (OCD) has been found to be similar in different countries and, within Western countries, between different ethnic groups, concern has been raised about the potential for differential equity of healthcare access for this patient group. Fernández de la Cruz *et al* (pp. 530–535) found that individuals with OCD from minority ethnic groups were underrepresented in the secondary and tertiary mental health services of one London catchment area. The authors found that the disparity was greater than that found for a comparison patient group with depression and they comment on the need for further research to focus on understanding the specific reasons for the identified inequality in service use. Also focused on differential patterns of healthcare access, Ran *et al* (pp. 495–500) have examined the outcomes of treated and never-treated patients with schizophrenia, in a study based in rural China. Over a 14-year follow-up period, those with schizophrenia who did not receive antipsychotic treatment were found to have poorer outcomes across both mental health and social domains. The authors argue that their findings challenge the notion that outcomes for those with serious mental illness are better in low- and middle-income countries.

In an editorial focused on the inequalities of physical health outcomes for those with mental illness, Shiers *et al* (pp. 471–473) call for more coordination between primary care, secondary care, and public health to address the poor physical health and mortality outcomes for individuals diagnosed with psychosis. The authors highlight a number of available resources designed to improve coordination of care with the aim of reducing the morbidity/mortality gap associated with psychosis. Another editorial in the *BJPsych* this month, by Taggart & Bailey (pp. 469–470), further highlights the excess mortality associated with serious mental illness in England by describing a mental-health-specific *Atlas of Variation*, which presents relevant life-course data by region. The authors call for inequity of healthcare access to be targeted in order to reduce the stubborn mortality gap.

Reported discrimination among people treated for depression is addressed in a cross-national study by Lasalvia *et al* (pp. 507–514), as part of the ASPEN/INDIGO international study. Of the 34 countries included in the study, people living in countries identified as having a very high Human Development Index (HDI) reported higher levels of discrimination than those in medium/low HDI countries. Examining the impact of individual and contextual factors in explaining differences and considering

the public awareness approaches taken in high-income countries to date, the authors propose a move away from stigma-reduction campaigns focused on the biological aetiology of mental illness to a focus on competence and inclusion.

Focusing on recovery

Three papers in the *BJPsych* this month address different aspects of recovery in mental health. Williams *et al* (pp. 551–555) evaluated the Questionnaire about the Process of Recovery (QPR) instrument and compared the 15- and 22-item version in two samples with longitudinal data. Both versions of the instrument were found to have satisfactory psychometric properties but the authors were able to recommend the shorter version as being slightly more robust and less burdensome. Using latent class growth analysis techniques, Hodgekins *et al* (pp. 536–543) identified three types of social recovery profile in a sample of individuals with first-episode psychosis followed up for 12 months, with 66% being classed in the low stable group. Such poor social recovery was predicted by male gender, ethnic minority status, young age at psychosis onset, increased negative symptoms and poor premorbid adjustment. From measurement and outcome prediction to development of an intervention to increase mental health team support for personal recovery, Slade *et al* (pp. 544–550) describe the process of evidence synthesis which led to development of REFOCUS, an empirically supported manualised intervention which will be evaluated in a multisite cluster randomised controlled trial. The authors highlight the active and free availability of developed materials from the project.

Criminal justice pathways and victimisation experiences

In the context of presenting to services with a first episode of psychosis, Bhui *et al* (pp. 523–529) found that criminal justice pathways were more common in violent presentations, where psychopathy levels were greater and where drug use was present. A first presentation to services via the criminal justice system was also more common among Black Caribbean and Black African patients to an extent not fully explained by other factors. The authors call for more qualitative hypothesis-generating and quantitative hypothesis-testing research to focus on understanding these findings.

Patients with severe mental illness (SMI) are known to be more likely to experience victimisation than those in the general population. In a Dutch study by de Mooij *et al* (pp. 515–522), almost a quarter of individuals with SMI reported violent victimisation compared with almost 10% of controls. Victims with SMI were more likely to be assaulted by someone known to them and, within the patient group, in-patients were the most likely to report victimisation.