

# BJPpsych

The British Journal of Psychiatry

**Deinstitutionalised patients, homelessness and imprisonment: systematic review**

Petr Winkler *et al*

**An exploratory randomised trial of a simple, brief psychological intervention to reduce subsequent suicidal ideation and behaviour in patients admitted to hospital for self-harm**

Christopher Armitage *et al*

**Suicides during pregnancy and 1 year postpartum in Sweden, 1980–2007**

Annika Esscher *et al*

**Assisted and unassisted suicide in men and women: longitudinal study of the Swiss population**

Nicole Steck *et al*





# BJPsych

## Contents

- A17 Editorial Board  
A19 Highlights of this issue

### Editorials

- 409 **Is sleep disruption a trigger for postpartum psychosis?**  
K. J. S. Lewis, R. G. Foster and I. R. Jones

- 411 **Mrs Noone – poems by doctors**  
Rowena Warwick

- 412 **Deinstitutionalisation does not increase imprisonment or homelessness**

T. T. Salisbury and G. Thornicroft

- 414 **Evolving eating disorder psychopathology: conceptualising muscularity-oriented disordered eating**

S. B. Murray, S. Griffiths and J. M. Mond

### Analysis

- 416 **Practising evidence-based medicine in an era of high placebo response: number needed to treat reconsidered**  
S. P. Roose, B. R. Rutherford, M. M. Wall and M. E. Thase

### Review articles

- 421 **Deinstitutionalised patients, homelessness and imprisonment: systematic review**  
P. Winkler, B. Barrett, P. McCrone, L. Cséry, M. Janoušková and C. Höschl

- 428 **The Anatomy of Melancholy by Robert Burton – reflection**  
German E. Berrios

- 429 **Behavioural and psychological symptoms in dementia and the challenges for family carers: systematic review**  
A. Feast, M. Orrell, G. Charlesworth, N. Melunsky, F. Poland and E. Moniz-Cook

### Papers

- 435 **Urban–rural differences in incidence rates of psychiatric disorders in Denmark**  
E. Vassos, E. Agerbo, O. Mors and C. B. Pedersen
- 441 **Role played by the SP4 gene in schizophrenia and major depressive disorder in the Han Chinese population**  
J. Chen, K. He, Q. Wang, Z. Li, J. Shen, T. Li, M. Wang, Z. Wen, W. Li, Y. Qiang, T. Wang, J. Ji, N. Wu, Z. Wang, B. Zhang, G. Feng, L. He, Y. Xu and Y. Shi
- 446 **Complement factor H and susceptibility to major depressive disorder in Han Chinese**  
C. Zhang, D.-F. Zhang, Z.-G. Wu, D.-H. Peng, J. Chen, J. Ni, W. Tang, L. Xu, Y.-G. Yao and Y.-R. Fang

- 452 **An 18th-century view of demonomania. 2: Vampirism – explanation – psychiatry in history**  
Fiona Subotsky

- 453 **Evaluation of ethnic disparities in detection of depression and anxiety in primary care during the maternal period: combined analysis of routine and cohort data**  
S. L. Prady, K. E. Pickett, E. S. Petherick, S. Gilbody, T. Croudace, D. Mason, T. A. Sheldon and J. Wright

- 462 **Suicides during pregnancy and 1 year postpartum in Sweden, 1980–2007**  
A. Esscher, B. Essén, E. Innala, F. C. Papadopoulos, A. Skalkidou, I. Sundström-Poromaa and U. Högberg

- 470 **An exploratory randomised trial of a simple, brief psychological intervention to reduce subsequent suicidal ideation and behaviour in patients admitted to hospital for self-harm**

C. J. Armitage, W. Abdul Rahim, R. Rowe and R. C. O'Connor

- 477 **Evaluation of benefit to patients of training mental health professionals in suicide guidelines: cluster randomised trial**  
D. P. de Beurs, M. H. de Groot, J. de Keijser, E. van Duijn, R. F. P. de Winter and A. J. F. M. Kerkhof

- 484 **Assisted and unassisted suicide in men and women: longitudinal study of the Swiss population**  
N. Steck, M. Egger and M. Zwahlen for the Swiss National Cohort

- 490 **Lovely young consultant charms my husband – poems by doctors**  
Kate Compston

- 491 **An evidence-based algorithm for the utility of FDG-PET for diagnosing Alzheimer's disease according to presence of medial temporal lobe atrophy**  
M. J. Firbank, J. Lloyd, D. Williams, R. Barber, S. J. Colloby, N. Barnett, K. Olsen, C. Davison, C. Donaldson, K. Herholz and J. T. O'Brien

### Short report

- 497 **Visual cortical excitability in dementia with Lewy bodies**  
J.-P. Taylor, M. Firbank and J. T. O'Brien

### Columns

- 499 Correspondence  
500 Correction  
500 Contents of *BJPsych Advances*  
501 Book reviews  
502 Contents of the *American Journal of Psychiatry*  
503 Kaleidoscope  
505 From the Editor's desk

### Cover picture

*Pain is a Thing with Feathers* (2015).  
Sophia Platts-Palmer (b.1991)

Sophia Platts-Palmer is a London-born illustrator living and working in Glasgow. Her work draws on her degree background in English literature and history of art, and is a hybrid of pictorial references to classicism and postmodern narrative themes. Much of her work addresses her own significant experiences with her mental health, womanhood and generational identity with a focus on the plurality of self, fluidity of gender, fractured identities and the comparisons between the female form and processes and phenomena of the natural world. Sophia has recently graduated with a masters in Curatorial Practice and Contemporary Art from the Glasgow School of Art.

*Pain is a Thing with Feathers* takes its name from Emily Dickinson's poem "Hope" is the Thing with Feathers' and is a monstrous representation of the artist's experience with fibromyalgia and the daily battle against pain and fatigue that the condition brings.

Twitter: splattspalmer

Website: [www.splattspalmer.com](http://www.splattspalmer.com)

We are always looking for interesting and visually appealing images for the cover of the *Journal* and would welcome suggestions or pictures, which should be sent to Dr Allan Beveridge, British Journal of Psychiatry, 21 Prescott Street, London E1 8BB, UK or [bjp@rcpsych.ac.uk](mailto:bjp@rcpsych.ac.uk).





## Highlights of this issue

By Derek K. Tracy

### Challenging preconceptions

Coming at the tail end of the era of the asylum, the Penrose (or hydraulic) hypothesis posited that deinstitutionalisation of long-term residential patients would lead to a rise in prison incarceration, and others have argued about further adverse sequelae such as greater rates of homelessness. The topic has a contemporary flavour: while we no longer have asylums, we live in a time of financial constraint and ever-tighter in-patient bed numbers. Winkler and colleagues (pp. 421–428) systematically review the issue and refute the core hypothesis, with no correlation identified between discharge after long-term care and subsequent homelessness and/or imprisonment. They note that the arguments that had supported the concept were often subjective editorials or ecological studies that could not disentangle concomitant confounders such as changes in globalisation, alterations to traditional support networks, and operational differences in mental health services. Efficacious public investment is the proposed 'hydraulic', not deinstitutionalisation; Salisbury & Thornicroft's editorial (pp. 412–413) advocates that the debate now needs to move to looking at the *optimal* in-patient/community ratio in differing settings.

Murray *et al* (pp. 414–415) note how much work in eating disorders focuses on female patients, meaning that professionals can be less sighted on men exhibiting such difficulties, not least as their distinct presentation can elude classification systems. It is a concern, especially as there is evidence of a growth in prevalence (in both genders), and the finding that men are often less likely to seek treatment for such behaviours. Some body dissatisfaction may fit with wider cultural trends of male 'action figures' with muscularity greater than that seen in the biggest of bodybuilders. Protein 'bulking', fat/carbohydrate 'cutting', and anabolic steroid use are argued to parallel the more typically female purging behaviour. Murray and colleagues propose that the transdiagnostic core is overvaluation of shape and weight, with various control behaviours secondary to these; further, they call for future diagnostic criteria to recognise this.

### Dementia: causes, detection, and living with illness

Dementia with Lewy bodies (DLB) accounts for 10–15% of degenerative dementias in older people. It is notable for its visual hallucinations, but it has proven difficult to elucidate their aetiology. Taylor and colleagues (pp. 497–498) compared transcranial magnetic stimulation (TMS) and functional magnetic resonance imaging (fMRI) data, finding a loss of inhibitory drive in the visual cortex in DLB that they suggest may drive the pathophysiology. Neuroimaging is used to diagnose dementia, with computed tomography (CT) detected medial temporal lobe atrophy a cardinal sign of Alzheimer's disease, delineating it from DLB. Temporoparietal hypometabolism, as measured by fluorodeoxyglucose positron emission tomography (FDG-PET), has also been proposed as a differentiator of these conditions but it is unclear how much value this test adds to the more standard (and cheaper and quicker) repertoire. Firbank *et al* (pp. 491–496) compared CT and FDG-PET across a sample that included individuals with Alzheimer's disease and DLB; they found that the novel investigation did add clinical utility, but only

in those with less (or no) atrophy, and was of limited use when losses in the medial temporal lobe were more severe.

Living with dementia is a profound burden for individuals and those around them. Their behaviour and psychological symptoms (BPSD), which afflict about 90%, are commonly a particular distress to family carers. Feast *et al* (pp. 429–434) review the BPSD factors that impact most upon the individual–carer dyad. Changes in communication and relationships led to family carers feeling bereft, while transgressions against social norms was the other factor identified as fundamental to their belief that their loved one had lost their identity to illness. Carers' acceptance of, and responses to, BPSD can vary considerably, and such factors have the power to alter the trajectory of these symptoms. The authors conclude that there is a significant unmet psychological need in family carers, who have lost an emotional bond, and have their own requirements for nurture, care and emotional security.

### Training and intervention in suicide

The incidence of suicide in the year following giving birth is *lower* than in women who have not delivered, but it is nevertheless an area of importance, and one of the commonest causes of death in women in high-income countries during this period. Esscher and colleagues (pp. 462–469) analysed almost 30 years of Swedish national register data, and determined a suicide ratio of 3.7/100 000 live births, but with an odds ratio over three times greater in women born in low-income countries. Violent methods of death were considerably more common than normal, occurring in 87% of those who died before the end of the 6-month postpartum period. Of concern, antenatal documentation was inconsistent, and a considerable number had no psychiatric care plan.

The complex issue of *assisted* suicide is addressed by Steck *et al* (pp. 484–490); the study was undertaken in Switzerland, one of the few countries to allow this act (provided there is no self-interest in such assistance, and euthanasia is banned). The overall suicide rate in Switzerland is higher than most European countries; most choosing assisted acts are terminally ill, but it is also legal in capacious healthy individuals. Over 1300 assisted and 5700 unassisted suicides were analysed. The results showed that rates of assisted suicide were similar in men and women (bucking the typical gender differences seen in unassisted acts). There were no differences based upon religious affiliation, having children or living alone; however, tertiary education was positively associated with assisted, and negatively associated with unassisted, suicide.

This month's Kaleidoscope (pp. 503–504) has fascinating data that the opioid partial agonist buprenorphine can acutely reduce suicidal thinking – the hypothesis being the anaesthetising of mental pain – but what about psychosocial inputs? There are some conflicting data from two papers: de Beurs *et al* (pp. 477–483) found no benefit from training professionals, via eLearning, in adhering to suicide prevention guidelines, while Armitage *et al* (pp. 470–476) showed that an 'implementation intentions' programme produced significant benefit. Suicide prevention training has been shown to improve the knowledge, skills and attitudes of practitioners. It is therefore a challenge understanding why training in the application of professional practice guidelines (and putatively therein the delivery of optimal evidence-based care) did not improve 3-month outcomes – in terms of suicide ideation, non-fatal attempts or treatment satisfaction – in this first randomised controlled trial on the topic. Notably, the control group also showed improvement, and the authors considered their sample size was insufficient to detect additional training benefits. The positive outcomes of the 'implementation intentions' programme, which links triggers for self-harm with coping skills, may suggest more focused interventions are necessary.